Calculus 1
Workbook Solutions

Modifying functions

krista king
MATH

COMBINATIONS OF FUNCTIONS

1. Find $(f+g)(x)$.

$$f(x) = 2x^2 - x + 5$$

$$g(x) = x^2 + 4x - 7$$

Solution:

The combination $(f+g)(x)$ is the same as $f(x) + g(x)$, so we need to add the equations together.

$$(f+g)(x) = f(x) + g(x)$$

$$(f+g)(x) = 2x^2 - x + 5 + x^2 + 4x - 7$$

Group like terms together and combine.

$$(f+g)(x) = 2x^2 + x^2 - x + 4x + 5 - 7$$

$$(f+g)(x) = 3x^2 + 3x - 2$$

2. Find $(f-g)(x)$.

$$f(x) = 4x^2 - 2$$

$$g(x) = 3x^2 - 5x$$

© Krista King Math

Solution:

The combination $(f-g)(x)$ is the same as $f(x)-g(x)$, so we need to find the difference.

$$(f-g)(x)=f(x)-g(x)$$

$$(f-g)(x)=4x^2-2-(3x^2-5x)$$

$$(f-g)(x)=4x^2-2-3x^2+5x$$

Group like terms together and combine.

$$(f-g)(x)=4x^2-3x^2+5x-2$$

$$(f-g)(x)=x^2+5x+2$$

3. Find $(f-g)(x)$.

$$f(x)=x^2-3x+1$$

$$g(x)=2x-3$$

Solution:

The combination $(f-g)(x)$ is the same as $f(x)-g(x)$, so we need to find the difference.

© Krista King Math

$$(f-g)(x)=f(x)-g(x)$$

$$(f-g)(x)=x^2-3x+1-(2x-3)$$

$$(f-g)(x)=x^2-3x+1-2x+3$$

Group like terms together and combine.

$$(f-g)(x)=x^2-3x-2x+1+3$$

$$(f-g)(x)=x^2-5x+4$$

4. Find $(f\cdot g)(x)$.

$$f(x)=2x-3$$

$$g(x)=3x^2+2$$

Solution:

The combination $(f\cdot g)(x)$ is the same as $f(x)\cdot g(x)$, so we need to find the product.

$$(f\cdot g)(x)=f(x)\cdot g(x)$$

$$(f\cdot g)(x)=(2x-3)(3x^2+2)$$

Multiply using the FOIL method.

$$(f\cdot g)(x)=6x^3+4x-9x^2-6$$

© Krista King Math

$$(f \cdot g)(x) = 6x^3 - 9x^2 + 4x - 6$$

5. Find $(f \cdot g)(x)$.

$$f(x) = x - 3$$

$$g(x) = x + 4$$

Solution:

The combination $(f \cdot g)(x)$ is the same as $f(x) \cdot g(x)$, so we need to find the product.

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$(f \cdot g)(x) = (x - 3)(x + 4)$$

Multiply using the FOIL method.

$$(f \cdot g)(x) = x^2 + 4x - 3x - 12$$

$$(f \cdot g)(x) = x^2 + x - 12$$

6. Find $(f \div g)(x)$.

$$f(x) = x^2 + 6x$$

$$g(x) = x$$

© Krista King Math

Solution:

The combination $(f \div g)(x)$ is the same as $f(x)/g(x)$, so we need to find the quotient.

$$(f \div g)(x) = \frac{x^2 + 6x}{x}$$

$$(f \div g)(x) = \frac{x(x+6)}{x}$$

$$(f \div g)(x) = x + 6$$

7. Find $(g \div f)(x)$.

$$f(x) = x^2 + 6x$$

$$g(x) = x$$

Solution:

The combination $(g \div f)(x)$ is the same as $g(x)/f(x)$, so we need to find the quotient.

$$(g \div f)(x) = \frac{x}{x^2 + 6x}$$

$$(g \div f)(x) = \frac{x}{x(x+6)}$$

© Krista King Math

$$(g \div f)(x) = \frac{1}{x+6}$$

© Krista King Math

COMPOSITE FUNCTIONS

1. Find the composite function $(f \circ g)(x)$.

$$f(x) = \sqrt{2x - 1}$$

$$g(x) = 3x^2$$

Solution:

When we take the composite $(f \circ g)(x)$, we plug $g(x)$ into $f(x)$.

$$(f \circ g)(x) = f(g(x)) = \sqrt{2(3x^2) - 1}$$

$$(f \circ g)(x) = f(g(x)) = \sqrt{6x^2 - 1}$$

2. Find the composite function $(g \circ f)(x)$.

$$f(x) = \sqrt{2x - 1}$$

$$g(x) = 3x^2$$

Solution:

When we take the composite $(f \circ g)(x)$, we plug $f(x)$ into $g(x)$.

© Krista King Math

$$(g \circ f)(x) = g(f(x)) = 3(\sqrt{2x-1})^2$$

$$(g \circ f)(x) = g(f(x)) = 3(2x-1)$$

$$(g \circ f)(x) = g(f(x)) = 6x-3$$

3. Find the composite function $f(g(x))$.

$$f(x) = x^2 - 4x + 3$$

$$g(x) = 2x + 1$$

Solution:

When we take the composite $f(g(x))$, we plug $g(x)$ into $f(x)$.

$$f(g(x)) = (2x+1)^2 - 4(2x+1) + 3$$

$$f(g(x)) = (2x+1)(2x+1) - 8x - 4 + 3$$

$$f(g(x)) = 4x^2 + 2x + 2x + 1 - 8x - 1$$

Group and combine like terms.

$$f(g(x)) = 4x^2 + 2x + 2x - 8x + 1 - 1$$

$$f(g(x)) = 4x^2 + 4x - 8x$$

$$f(g(x)) = 4x^2 - 4x$$

© Krista King Math

4. Find the composite function $g(f(x))$.

$$f(x) = x^2 - 4x + 3$$

$$g(x) = 2x + 1$$

Solution:

When we take the composite $g(f(x))$, we plug $f(x)$ into $g(x)$.

$$g(f(x)) = 2(x^2 - 4x + 3) + 1$$

$$g(f(x)) = 2x^2 - 8x + 6 + 1$$

$$g(f(x)) = 2x^2 - 8x + 7$$

5. Find the composite function $(g \circ h)(x)$.

$$g(x) = \frac{8}{x^3}$$

$$h(x) = \sqrt[3]{x + 4}$$

Solution:

When we take the composite $(g \circ h)(x)$, we plug $h(x)$ into $g(x)$.

$$(g \circ h)(x) = g(h(x)) = \frac{8}{(\sqrt[3]{x + 4})^3}$$

© Krista King Math

$$(g \circ h)(x) = g(h(x)) = \frac{8}{x+4}$$

6. Find the composite function $(h \circ g)(x)$.

$$g(x) = \frac{8}{x^3}$$

$$h(x) = \sqrt[3]{x+4}$$

Solution:

When we take the composite $(h \circ g)(x)$, we plug $g(x)$ into $h(x)$.

$$(h \circ g)(x) = h(g(x)) = \sqrt[3]{\frac{8}{x^3} + 4}$$

$$(h \circ g)(x) = h(g(x)) = \sqrt[3]{\frac{8}{x^3} + 4\frac{x^3}{x^3}}$$

$$(h \circ g)(x) = h(g(x)) = \sqrt[3]{\frac{8+4x^3}{x^3}}$$

Take the root of the numerator and denominator separately.

$$(h \circ g)(x) = h(g(x)) = \frac{\sqrt[3]{8+4x^3}}{\sqrt[3]{x^3}}$$

© Krista King Math

$$(h \circ g)(x) = h(g(x)) = \frac{\sqrt[3]{8 + 4x^3}}{x}$$

7. Find the composite function $g(h(x))$.

$$g(x) = \frac{1}{x}$$

$$h(x) = 3x^2 - x$$

Solution:

When we take the composite $g(h(x))$, we plug $h(x)$ into $g(x)$.

$$g(h(x)) = \frac{1}{3x^2 - x}$$

8. Find the composite function $h(g(x))$.

$$g(x) = \frac{1}{x}$$

$$h(x) = 3x^2 - x$$

Solution:

When we take the composite $h(g(x))$, we plug $g(x)$ into $h(x)$.

© Krista King Math

$$h(g(x)) = 3\left(\frac{1}{x}\right)^2 - \frac{1}{x}$$

$$h(g(x)) = \frac{3}{x^2} - \frac{1}{x}$$

Find a common denominator to combine the fractions.

$$h(g(x)) = \frac{3}{x^2} - \frac{1}{x}\left(\frac{x}{x}\right)$$

$$h(g(x)) = \frac{3}{x^2} - \frac{x}{x^2}$$

$$h(g(x)) = \frac{3-x}{x^2}$$

© Krista King Math

COMPOSITE FUNCTIONS, DOMAIN

1. What is the domain of $f \circ g$?

$$f(x) = x^2 - 2$$

$$g(x) = \sqrt{x + 3}$$

Solution:

Find the domain of $g(x)$. Remember that you can't take the square root of negative numbers, since negative roots can't be defined by real numbers, so the expression inside the root must be positive or equal to 0.

$$x + 3 \geq 0$$

$$x \geq -3$$

Now find $f \circ g$.

$$f \circ g = (\sqrt{x + 3})^2 - 2$$

$$f \circ g = x + 3 - 2$$

$$f \circ g = x + 1$$

We need to consider the domain of this composite function. The composite is a simple binomial with no domain restrictions. There's only the domain restriction that we found for $g(x)$, but all restrictions of $g(x)$

© Krista King Math

need to be included in the domain for $f(g(x))$. Therefore, the domain of $f(g(x))$ is $x \geq -3$.

2. What is the domain of $f \circ g$?

$$f(x) = \frac{1}{x}$$

$$g(x) = x + 5$$

Solution:

Find the domain of $g(x)$. In this case, $g(x)$ is a simple binomial with no domain restrictions.

Now find $f \circ g$.

$$f \circ g = \frac{1}{x + 5}$$

We need to consider the domain of this composite function. Remember that we can't divide by 0, which means the denominator of a fraction can never equal 0.

$$x + 5 \neq 0$$

$$x \neq -5$$

There were no restrictions on the domain of $g(x)$, so the domain of $f(g(x))$ is just $x \neq -5$.

© Krista King Math

3. What is the domain of $f \circ g$?

$$f(x) = \frac{2}{x-1}$$

$$g(x) = \sqrt{x-4}$$

Solution:

Find the domain of $g(x)$. Remember that you can't take the square root of negative numbers, since negative roots can't be defined by real numbers, so the expression inside the root must be positive or equal to 0.

$$x - 4 \geq 0$$

$$x \geq 4$$

Now find $f \circ g$.

$$f \circ g = \frac{2}{\sqrt{x-4}-1}$$

We need to consider the domain of this composite function. Remember that we can't divide by 0, which means the denominator of a fraction can never equal 0.

$$\sqrt{x-4} - 1 \neq 0$$

$$\sqrt{x-4} \neq 1$$

© Krista King Math

$$x - 4 \neq 1$$

$$x \neq 5$$

The domain of the composite has to account for domain restrictions on $g(x)$ and the composite function $f \circ g$ itself, so the domain of $f(g(x))$ is $x \geq 4,\ x \neq 5$.

4. What is the domain of $f \circ g$?

$$f(x) = \frac{1}{x} + 4$$

$$g(x) = \frac{3}{2x - 7}$$

Solution:

Find the domain of $g(x)$. Remember that we can't divide by 0, which means the denominator of a fraction can never equal 0.

$$2x - 7 \neq 0$$

$$2x \neq 7$$

$$x \neq \frac{7}{2}$$

Now find $f \circ g$.

© Krista King Math

$$f \circ g = \frac{1}{\frac{3}{2x-7}} + 4$$

$$f \circ g = \frac{2x-7}{3} + 4$$

$$f \circ g = \frac{2x-7}{3} + \frac{12}{3}$$

$$f \circ g = \frac{2x-7+12}{3}$$

$$f \circ g = \frac{2x+5}{3}$$

We need to consider the domain of this composite function. But there are no domain restrictions on $f \circ g$, so the only restriction we need to consider is the one on $g(x)$. The domain of $f(g(x))$ is $x \neq 7/2$.

5. What is the domain of $f \circ g$?

$$f(x) = \frac{2}{x-3}$$

$$g(x) = \frac{4}{x+2}$$

Solution:

© Krista King Math

Find the domain of $g(x)$. Remember that we can't divide by 0, which means the denominator of a fraction can never equal 0.

$$x + 2 \neq 0$$

$$x \neq -2$$

Now find $f \circ g$.

$$f \circ g = \frac{2}{\frac{4}{x+2} - 3}$$

$$f \circ g = \frac{2}{\frac{4}{x+2} - 3\left(\frac{x+2}{x+2}\right)}$$

$$f \circ g = \frac{2}{\frac{4}{x+2} - \frac{3x+6}{x+2}}$$

Combine fractions within the denominator.

$$f \circ g = \frac{2}{\frac{4-3x-6}{x+2}}$$

$$f \circ g = \frac{2}{\frac{-3x-2}{x+2}}$$

$$f \circ g = \frac{2(x+2)}{-3x-2}$$

$$f \circ g = \frac{2x+4}{-3x-2}$$

© Krista King Math

$$f \circ g = -\frac{2x+4}{3x+2}$$

We need to consider the domain of this composite function. Remember that we can't divide by 0, which means the denominator of a fraction can never equal 0.

$$3x + 2 \neq 0$$

$$3x \neq -2$$

$$x \neq -\frac{2}{3}$$

Putting this restriction together with the one we found for $g(x)$, we can say that the domain of $f(g(x))$ is $x \neq -2, \ -2/3$.

6. What is the domain of $f \circ g$?

$$f(x) = \frac{1}{x^2 - 3}$$

$$g(x) = \sqrt{x - 1}$$

Solution:

Find the domain of $g(x)$. Remember that you can't take the square root of negative numbers, since negative roots can't be defined by real numbers, so the expression inside the root must be positive or equal to 0.

© Krista King Math

$$x - 1 \geq 0$$

$$x \geq 1$$

Now find $f \circ g$.

$$f \circ g = \frac{1}{(\sqrt{x-1})^2 - 3}$$

$$f \circ g = \frac{1}{x - 1 - 3}$$

$$f \circ g = \frac{1}{x - 4}$$

We need to consider the domain of this composite function. Remember that we can't divide by 0, which means the denominator of a fraction can never equal 0.

$$x - 4 \neq 0$$

$$x \neq 4$$

Putting this restriction together with the one we found for $g(x)$, we can say that the domain of $f(g(x))$ is $x \geq 1$, $x \neq 4$.

7. What is the domain of $f \circ g$?

$$f(x) = 2x^2 - x + 1$$

$$g(x) = x - 3$$

© Krista King Math

Solution:

Find the domain of $g(x)$. In this case $g(x)$ is a simple binomial with no domain restrictions.

Now find $f \circ g$.

$$f \circ g = 2(x-3)^2 - (x-3) + 1$$

$$f \circ g = 2(x^2 - 6x + 9) - x + 3 + 1$$

$$f \circ g = 2x^2 - 12x + 18 - x + 4$$

$$f \circ g = 2x^2 - 13x + 22$$

We need to consider the domain of this composite function. But there are no domain restrictions on $f \circ g$. And, since there are also no domain restrictions for $g(x)$, the domain of $f(g(x))$ is all real numbers.

■ 8. What is the domain of $f \circ g$?

$$f(x) = x^2 + 4x - 10$$

$$g(x) = x + 6$$

Solution:

© Krista King Math

Find the domain of $g(x)$. In this case $g(x)$ is a simple binomial with no domain restrictions.

Now find $f \circ g$.

$$f \circ g = (x+6)^2 + 4(x+6) - 10$$

$$f \circ g = x^2 + 12x + 36 + 4x + 24 - 10$$

$$f \circ g = x^2 + 16x + 50$$

There are no domain restrictions for $f \circ g$. And, since there are also no domain restrictions for $g(x)$, the domain of $f(g(x))$ is all real numbers.

© Krista King Math

FOR THE LOVE OF MATH

WWW.KRISTAKINGMATH.COM